



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

APRIL 17.

The President, SAMUEL G. DIXON, M.D., in the Chair.

One hundred persons present.

A paper entitled "A Review of the Physæ of Northeastern Illinois," by Frank C. Baker, was presented for publication.

The death of Charles E. Smith, a member, on the 15th inst., was announced.

Respiratory Quotient and Loss in Volume of Expired Air.—DR. HENRY C. CHAPMAN called attention to the fact that the so-called "respiratory quotient," as originally defined by Pflüger¹ is the ratio of the weight of the oxygen absorbed during inspiration to the weight of the oxygen in the carbon dioxide exhaled during expiration. Inasmuch, however, as the volume of carbon dioxide formed is equal to the volume of oxygen entering into its formation, the "respiratory quotient" is usually defined as being the ratio of the volume of oxygen absorbed to the volume of carbon dioxide exhaled. Thus, for example, on the supposition that 100 litres of air be inspired, that 4.78 litres = 6.834 gr. of oxygen be absorbed, and that 4.34 litres = 6.205 gr. of oxygen be exhaled, the respiratory quotient will be

$$\frac{\text{Oxygen } 6.205 \text{ gr.}}{\text{Oxygen } 6.834} = \frac{4.34 \text{ lit.}}{4.78} = 0.907$$

or

$$\frac{\text{Carbon dioxide } 4.34 \text{ lit.}}{\text{Oxygen } 4.78} = 0.907$$

It will be observed that in making use of the expression "respiratory quotient" in the sense used by Pflüger, it is indifferent whether the oxygen absorbed and that exhaled in the carbon dioxide be estimated in grammes or litres, but that in the second case the ratio will be only 0.907, when the ratio of volumes are compared. It is also quite obvious, though it appears to have hitherto escaped the attention of physiologists, that on the supposition that the "respiratory quotient" = 0.907, the loss in the volume of the expired air, as compared with that inspired, must be less than the one-fiftieth of a volume, as usually stated, and as originally determined by Despretz,² rather the one-two hundred and fiftieth

¹ Pflüger's *Archiv*, XIV, 1877, S. 472.

² *Annales de chimie et de physique*, XXVI, p. 337.

of a volume, since 4.78 minus $4.34 = 0.4$ the loss, and $\frac{1.00}{0.4} = 250$. It may be mentioned in this connection that the loss in the volume of the expired air is due to the fact that all of the oxygen absorbed does not reappear in the carbon dioxide exhaled, part of it forming in the economy other combinations, such as water, and to some extent also, sulphuric and phosphoric acids, etc. In order to avoid misunderstanding, it should be stated that as a matter of fact the volume of the expired air is greater than that inspired, on account of the usually higher temperature of the former. When, however, the volumes of the inspired and expired air are reduced to standard temperature and pressure, then the volume of the expired air will be found to be less than that inspired, the loss being about that just stated, varying according to the respiratory quotient, the latter depending in turn upon the diet.

MR. D. SHEPHERD HOLMAN made an illustrated communication on sound-waves. (No abstract.)

APRIL 24.

The President, SAMUEL G. DIXON, M.D., in the Chair.

Twenty-seven persons present.

PHILIP P. CALVERT, Ph.D., was appointed a member of the Committee on Publications, to fill the vacancy caused by the death of Charles E. Smith.

The deaths of William Camac, M.D., a member, and of Prof. Alphonse Milne Edwards, a correspondent, were announced.

Walter T. Taggart and Milo G. Miller, M.D., were elected members.

The following were ordered to be printed: